

Access DB# _____

Scientific and Technical Information Center

If more than one search is submitted, please prioritize searches in order of need.



Title of Invention: _____

Inventors (please provide full names): _____

Earliest Priority Filing Date: _____

**For Sequence Searches Only* Please include all pertinent information (parent, child, divisional, or issued patent numbers) along with the appropriate serial number.*

2 Searches



1) $Q = \text{NH}_2$, comp'd must be mult. component

2) Q = c1ccccc1C(=O)Cn $n = 1-6$. Compd cannot be multicomponent

STAFF USE ONLY

Searcher: N. J. J.

Searcher Phone #: _____

Searcher Location:

Date Searcher Picked Up: _____

Date Completed: 6/18

Searcher Prep & Review Time:

Clerical Prep Time: _____

Online Time: 27

Type of Search

NA Sequence (#)_____

AA Sequence (#)_____

Structure (#)

Bibliographic

Litigation _____

Fulltext

Patent Family _____

Other _____

Vendors and cost where applicable

STN 130.50

Dialog _____

Questel/Orbit _____

Dr. Link _____

Lexis/Nexis



Sequence Systems _____

WWW/Internet _____

Other (specify) _____

Burch
10/006579

=> d l8 que stat;d 1-3 ide cbib abs
L6 STR



VAR G1=29/26
REP G2=(1-6) C
VAR G3=NH2/38
NODE ATTRIBUTES:
DEFAULT MLEVEL IS ATOM
DEFAULT ECLEVEL IS LIMITED

GRAPH ATTRIBUTES:
RING(S) ARE ISOLATED OR EMBEDDED
NUMBER OF NODES IS 39

STEREO ATTRIBUTES: NONE
L8 3 SEA FILE=REGISTRY SSS FUL L6


100.0% PROCESSED 85 ITERATIONS
SEARCH TIME: 00.00.03

3 ANSWERS

L8 ANSWER 1 OF 3 REGISTRY COPYRIGHT 2002 ACS
RN 141194-86-7 REGISTRY
CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid,
7-amino-8-oxo-3-(tetrahydro-2-furanyl)-, (4-methoxyphenyl)methyl ester,
[6R-(6.alpha.,7.beta.)]- (9CI) (CA INDEX NAME)
FS STEREOSEARCH
MF C19 H22 N2 O5 S
SR CA
LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.

Searched by: Mary Hale 308-4258 CM-1 1E01



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1967 TO DATE)
1 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L8 ANSWER 2 OF 3 REGISTRY COPYRIGHT 2002 ACS

RN 141061-23-6 REGISTRY

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-amino-8-oxo-3-(tetrahydro-2-furanyl)-, (4-methoxyphenyl)methyl ester, [6R-[3(R*),6.alpha.,7.beta.]]- (9CI) (CA INDEX NAME)


FS STEREOSEARCH

MF C19 H22 N2 O5 S

SR CA

LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1967 TO DATE)

1 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L8 ANSWER 3 OF 3 REGISTRY COPYRIGHT 2002 ACS

RN 141061-22-5 REGISTRY

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-amino-8-oxo-3-(tetrahydro-2-furanyl)-, (4-methoxyphenyl)methyl ester, [6R-[3(S*),6.alpha.,7.beta.]]- (9CI) (CA INDEX NAME)


FS STEREOSEARCH

MF C19 H22 N2 O5 S

SR CA

LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.



PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT

1 REFERENCES IN FILE CA (1967 TO DATE)
1 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

=> scr 2127

L9 SCREEN CREATED

=> search

ENTER LOGIC EXPRESSION, QUERY NAME, OR (END):l9

ENTER TYPE OF SEARCH (SSS), CSS, FAMILY, OR EXACT:.

ENTER SCOPE OF SEARCH (SAMPLE), FULL, RANGE, OR SUBSET:subset

ENTER SUBSET L# OR (END):l8

ENTER SUBSET SEARCH SCOPE - SAMPLE, FULL, RANGE, OR (END):ful

FULL SUBSET SEARCH INITIATED 16:17:48

FULL SUBSET SCREEN SEARCH COMPLETED

0 ANSWERS

SEARCH TIME: 00.00.01


L10 0 SEA SUB=L8 SSS FUL L9

=> d l13 que stat;d 1-13 ide cbib abs

L9 SCR 2127

L11 STR

Searched by: Mary Hale 308-4258 CM-1 1E01



NODE ATTRIBUTES:
 DEFAULT MLEVEL IS ATOM
 DEFAULT ECLEVEL IS LIMITED


GRAPH ATTRIBUTES:
 RING(S) ARE ISOLATED OR EMBEDDED
 NUMBER OF NODES IS 17

STEREO ATTRIBUTES: NONE
 L13 13 SEA FILE=REGISTRY SSS FUL L11 AND L9

100.0% PROCESSED 53 ITERATIONS 13 ANSWERS
 SEARCH TIME: 00.00.01

L13 ANSWER 1 OF 13 REGISTRY COPYRIGHT 2002 ACS
 RN 395661-01-5 REGISTRY
 CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid,
 8-oxo-7-[(phenylacetyl)amino]-3-[(2S)-tetrahydro-2-furanyl]-, monosodium
 salt, (6R,7R)- (9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C19 H20 N2 O5 S . Na
 SR CA
 LC STN Files: CA, CAPLUS, USPATFULL

Absolute stereochemistry.



● Na

1 REFERENCES IN FILE CA (1967 TO DATE)
 1 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 136:151036 Process for the preparation of cephalosporin

Searched by: Mary Hale 308-4258 CM-1 1E01

compounds and their intermediates. Burton, George; Best, Desmond John; Gasson, Brian Charles; Osborne, Neal Frederick; Walker, Graham (Pfizer Inc., USA). Eur. Pat. Appl. EP 1178049 A1 20020206, 22 pp. DESIGNATED STATES: R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO. (English). CODEN: EPXXDW. APPLICATION: EP 2001-306325 20010723. PRIORITY: GB 2000-19124 20000803.

GI

* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT *

AB A process for prepg. cephalosporins I (R1 = H, OMe, formamido; R2 = acyl; CO2R3 = carboxy group or CO2- or readily removable carboxy protecting group; R4 = H, or up to four substituents from alkyl, alkenyl, alkynyl, alkoxy, halogen, amino, alkyl(acyl)amino, CO2R, CONR2, SO2NR2 (R = H, C1-6 alkyl), aryl, heterocycle, etc.; X = S, SO, SO2, O, CH2; m = 1-2; dotted lines indicate a 2- or 3-cephem system) was accomplished via the cyclization of II. Thus the 3-(R and S)-tetrahydrofuran-2-yl-2-em compds. III were prepd. and the S isomer was converted to the 3-(S)-tetrahydrofuran-2-yl-3-em III in several steps.


L13 ANSWER 2 OF 13 REGISTRY COPYRIGHT 2002 ACS
 RN 179238-43-8 REGISTRY
 CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid,
 7-amino-8-oxo-3-(tetrahydro-2-furanyl)-, (2,2-dimethyl-1-oxopropoxy)methyl
 ester, [6R-[3(S*),6.alpha.,7.beta.]]-, mono(4-methylbenzenesulfonate)
 (9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C17 H24 N2 O6 S . C7 H8 O3 S
 SR CA
 LC STN Files: CA, CAPLUS

CM 1

CRN 141072-36-8

CMF C17 H24 N2 O6 S


Absolute stereochemistry.



CM 2

CRN 104-15-4


CMF C7 H8 O3 S



1 REFERENCES IN FILE CA (1967 TO DATE)
1 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 125:114393 Process for the preparation of cephalosporins and analogs. Burton, George; Naylor, Antoinette (Pfizer Inc., USA). PCT Int. Appl. WO 9617847 A1 19960613, 29 pp. DESIGNATED STATES: W: JP, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1995-GB2783 19951129. PRIORITY: GB 1994-24847 19941209.

GI



AB Cephalosporins I [X = S, SO, SO₂, O, CH₂; R₁ = H, OMe, NHCHO; R₂ = acyl; R₃ = in vivo hydrolyzable ester group; R₄ = (un)substituted tetrahydrofuryl, tetrahydropyranyl] are prepd. by reaction of the corresponding carboxylic acid with R₃Y [Y = halide] in the presence of an aq. phase contg. a base and a phase transfer catalyst. Subsequent removal of protecting groups, conversion of groups X and R₂ and salt formation may be carried out. Thus, 4-methoxybenzyl (6R,7R)-7-phenylacetamido-3-[(S)-2-tetrahydrofuryl]cephem-4-carboxylate was treated with Me₃CCO₂CH₂I, followed by deacylation and reacylation to give pivaloyloxymethyl (6R,7R)-7-[2-(2-amino-4-thiazolyl)-2-(Z)-methoxyiminoacetamido]-3-[(S)-2-tetrahydrofuryl]cephem-4-carboxylate.

L13 ANSWER 3 OF 13 REGISTRY COPYRIGHT 2002 ACS

RN 141195-79-1 REGISTRY

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[[amino(4-hydroxyphenyl)acetyl]amino]-8-oxo-3-(tetrahydro-2-furanyl)-, monosodium salt, [6R-[6.alpha.,7.beta.(R*)]]- (9CI) (CA INDEX NAME)


FS STEREOSEARCH

MF C19 H21 N3 O6 S . Na

SR CA

LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.



● Na

1 REFERENCES IN FILE CA (1967 TO DATE)
1 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L13 ANSWER 4 OF 13 REGISTRY COPYRIGHT 2002 ACS

RN 141195-78-0 REGISTRY

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[[[(2-amino-4-thiazolyl)(methoxyimino)acetyl]amino]-8-oxo-3-(tetrahydro-2-furanyl)-, monosodium salt, [6R-[3(R*),6.alpha.,7.beta.(Z)]]- (9CI) (CA INDEX NAME)

FS STEREOSEARCH


MF C17 H19 N5 O6 S2 . Na

SR CA

LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.

Double bond geometry as shown.




● Na


2 REFERENCES IN FILE CA (1967 TO DATE)
2 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 126:144046 Beta-lactam preparation. Harris, Michael Anthony; Saunders, Richard Neville (Pfizer Limited, UK). Brit. UK Pat. Appl. GB 2300856 A1 19961120, 15 pp. (English). CODEN: BAXXDU. APPLICATION: GB 1995-10126 19950516.

GI



I



II

AB Title compds. I [R = substituent; R1 = H, OMe, NHCHO; R2 = acyl; CO2R3 = CO2H, CO2-; R3 = protecting group; X = S, SO, SO2, O, CH2] are prepd. by base-induced cyclization of an azetidinone II [R4 = alkyl, aryl]. II are prepd. from the halide and P(OR4)3. Thus, 4-methoxybenzyl (2RS)-2-hydroxy-2-[(3R)(4R)-3-phenylacetamido-4-[(RS)-2-tetrahydrofuryl]carbonylmethylthio]azetidin-2-on-1-ylacetate was converted to the chloride and then to the phosphonate which was cyclized with NaH in PhMe to give 50% I [R = (RS)-2-tetrahydrofuryl, R1 = H, R2 = PhCH2CO, R3 = 4-MeC6H4CH2].

REFERENCE 2: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

Searched by: Mary Hale 308-4258 CM-1 1E01

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L13 ANSWER 5 OF 13 REGISTRY COPYRIGHT 2002 ACS

RN 141195-77-9 REGISTRY

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[[[(2Z)-(2-amino-4-thiazolyl)(methoxyimino)acetyl]amino]-8-oxo-3-[(2S)-tetrahydro-2-furanyl]-, monosodium salt, (6R,7R)- (9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[[[(2-amino-4-thiazolyl)(methoxyimino)acetyl]amino]-8-oxo-3-(tetrahydro-2-furanyl)-, monosodium salt, [6R-[3(S*),6.alpha.,7.beta.(Z)]]-

OTHER NAMES:

CN Cefovecin sodium

CN UK 287074-02

FS STEREOSEARCH

MF C17 H19 N5 O6 S2 . Na


SR CA

LC STN Files: CA, CAPLUS, TOXCENTER

CRN (234096-34-5)

Absolute stereochemistry.

Double bond geometry as shown.



● Na



2 REFERENCES IN FILE CA (1967 TO DATE)

2 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 126:144046 Beta-lactam preparation. Harris, Michael Anthony; Saunders, Richard Neville (Pfizer Limited, UK). Brit. UK Pat. Appl. GB 2300856 A1 19961120, 15 pp. (English). CODEN: BAXXDU. APPLICATION: GB 1995-10126 19950516.

GI

Searched by: Mary Hale 308-4258 CM-1 1E01



AB Title compds. I [R = substituent; R1 = H, OMe, NHCHO; R2 = acyl; CO2R3 = CO2H, CO2-; R3 = protecting group; X = S, SO, SO2, O, CH2] are prep'd. by base-induced cyclization of an azetidinone II [R4 = alkyl, aryl]. II are prep'd. from the halide and P(OR4)3. Thus, 4-methoxybenzyl (2RS)-2-hydroxy-2-[(3R)(4R)-3-phenylacetamido-4-[(RS)-2-tetrahydrofuryl]carbonylmethylthio]azetidin-2-on-1-ylacetate was converted to the chloride and then to the phosphonate which was cyclized with NaH in PhMe to give 50% I [R = (RS)-2-tetrahydrofuryl, R1 = H, R2 = PhCH2CO, R3 = 4-MeC6H4CH2].

REFERENCE 2: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prep'd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L13 ANSWER 6 OF 13 REGISTRY COPYRIGHT 2002 ACS

RN 141096-61-9 REGISTRY

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[[[(2-amino-4-thiazolyl)[(carboxymethoxy)imino]acetyl]amino]-8-oxo-3-(tetrahydro-2-furanyl)-, disodium salt, [6R-[6.alpha.,7.beta.(Z)]]- (9CI) (CA INDEX NAME)


FS STEREOSEARCH

MF C18 H19 N5 O8 S2 . 2 Na

SR CA

LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.
Double bond geometry as shown.



REFERENCE 1: 126:144046 Beta-lactam preparation. Harris, Michael Anthony; Saunders, Richard Neville (Pfizer Limited, UK). Brit. UK Pat. Appl. GB 2300856 A1 19961120, 15 pp. (English). CODEN: BAXXDU. APPLICATION: GB 1995-10126 19950516.

$$\begin{array}{c}
 \text{R}^2\text{NH} \\
 \diagup \\
 \text{C} \\
 \diagdown \\
 \text{O} \\
 \text{N} \\
 \diagup \\
 \text{C} \\
 \diagdown \\
 \text{CO}_2\text{R}^3
 \end{array}
 \begin{array}{c}
 \text{H} \\
 \vdots \\
 \text{X} - \text{COR} \\
 \text{P(O)(OR}^4\text{)}_2
 \end{array}
 \quad \text{II}$$


REFERENCE 2: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge,

carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2; SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L13 ANSWER 7 OF 13 REGISTRY COPYRIGHT 2002 ACS
 RN 141096-60-8 REGISTRY
 CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid,
 7-[[2-(2-amino-4-thiazolyl)-1-oxo-2-pentenyl]amino]-8-oxo-3-(tetrahydro-2-
 furanyl)-, monosodium salt, [6R-[3(S*),6.alpha.,7.beta.(Z)]]- (9CI) (CA
 INDEX NAME)
 FS STEREOSEARCH
 MF C19 H22 N4 O5 S2 . Na
 SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.
 Double bond geometry as shown.





● Na

2 REFERENCES IN FILE CA (1967 TO DATE)
 2 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 126:144046 Beta-lactam preparation. Harris, Michael Anthony; Saunders, Richard Neville (Pfizer Limited, UK). Brit. UK Pat. Appl. GB 2300856 A1 19961120, 15 pp. (English). CODEN: BAXXDU. APPLICATION: GB 1995-10126 19950516.

GI



AB Title compds. I [R = substituent; R1 = H, OMe, NHCHO; R2 = acyl; CO2R3 = CO2H, CO2-; R3 = protecting group; X = S, SO, SO2, O, CH2] are prep'd. by base-induced cyclization of an azetidinone II [R4 = alkyl, aryl]. II are prep'd. from the halide and P(OR4)3. Thus, 4-methoxybenzyl (2RS)-2-hydroxy-2-[(3R)(4R)-3-phenylacetamido-4-[(RS)-2-tetrahydrofuryl]carbonylmethylthio]azetidin-2-on-1-ylacetate was converted to the chloride and then to the phosphonate which was cyclized with NaH in PhMe to give 50% I [R = (RS)-2-tetrahydrofuryl, R1 = H, R2 = PhCH2CO, R3 = 4-MeC6H4CH2].

REFERENCE 2: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prep'd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L13 ANSWER 8 OF 13 REGISTRY COPYRIGHT 2002 ACS

RN 141082-25-9 REGISTRY

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[[2-furanyl(methoxyimino)acetyl]amino]-8-oxo-3-(tetrahydro-2-furanyl)-, monosodium salt, [6R-[3(S*),6.alpha.,7.beta.(Z)]]- (9CI) (CA INDEX NAME)

FS STEREOSEARCH


MF C18 H19 N3 O7 S . Na

SR CA

LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.

Double bond geometry as shown.





● Na

2 REFERENCES IN FILE CA (1967 TO DATE)

2 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 126:144046 Beta-lactam preparation. Harris, Michael Anthony; Saunders, Richard Neville (Pfizer Limited, UK). Brit. UK Pat. Appl. GB 2300856 A1 19961120, 15 pp. (English). CODEN: BAXXDU. APPLICATION: GB 1995-10126 19950516.

GI



AB Title compds. I [R = substituent; R1 = H, OMe, NHCHO; R2 = acyl; CO2R3 = CO2H, CO2-; R3 = protecting group; X = S, SO, SO2, O, CH2] are prep'd. by base-induced cyclization of an azetidinone II [R4 = alkyl, aryl]. II are prep'd. from the halide and P(OR4)3. Thus, 4-methoxybenzyl (2RS)-2-hydroxy-2-[(3R)(4R)-3-phenylacetamido-4-[(RS)-2-tetrahydrofuryl]carbonylmethylthio]azetidin-2-on-1-ylacetate was converted to the chloride and then to the phosphonate which was cyclized with NaH in PhMe to give 50% I [R = (RS)-2-tetrahydrofuryl, R1 = H, R2 = PhCH2CO, R3 = 4-MeC6H4CH2].

REFERENCE 2: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.


GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = 1 to 4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n = 0-2; m = 1, 2) were prep'd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-

yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L13 ANSWER 9 OF 13 REGISTRY COPYRIGHT 2002 ACS
 RN 141082-24-8 REGISTRY
 CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[[[(2-amino-4-thiazolyl)(methoxyimino)acetyl]amino]-8-oxo-3-(tetrahydro-2-furanyl)-, 5,5-dioxide, monosodium salt, [6R-[3(S*),6.alpha.,7.beta.(Z)]]-(9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C17 H19 N5 O8 S2 . Na
 SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.
 Double bond geometry as shown.





● Na

2 REFERENCES IN FILE CA (1967 TO DATE)
 2 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 126:144046 Beta-lactam preparation. Harris, Michael Anthony; Saunders, Richard Neville (Pfizer Limited, UK). Brit. UK Pat. Appl. GB 2300856 A1 19961120, 15 pp. (English). CODEN: BAXXDU. APPLICATION: GB 1995-10126 19950516.

GI



Searched by: Mary Hale 308-4258 CM-1 1E01

AB Title compds. I [R = substituent; R1 = H, OMe, NHCHO; R2 = acyl; CO2R3 = CO2H, CO2-; R3 = protecting group; X = S, SO, SO2, O, CH2] are prepd. by base-induced cyclization of an azetidinone II [R4 = alkyl, aryl]. II are prepd. from the halide and P(OR4)3. Thus, 4-methoxybenzyl (2RS)-2-hydroxy-2-[(3R)(4R)-3-phenylacetamido-4-[(RS)-2-tetrahydrofuryl]carbonylmethylthio]azetidin-2-on-1-ylacetate was converted to the chloride and then to the phosphonate which was cyclized with NaH in PhMe to give 50% I [R = (RS)-2-tetrahydrofuryl, R1 = H, R2 = PhCH2CO, R3 = 4-MeC6H4CH2].

REFERENCE 2: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L13 ANSWER 10 OF 13 REGISTRY COPYRIGHT 2002 ACS

RN 141082-22-6 REGISTRY

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[[[(2-amino-4-thiazolyl)(methoxyimino)acetyl]amino]-8-oxo-3-(tetrahydro-2-furanyl)-, 5-oxide, monosodium salt, [5S-[3(R*),5.alpha.,6.beta.,7.alpha.(Z)]]-(9CI) (CA INDEX NAME)

FS STEREOSEARCH


MF C17 H19 N5 O7 S2 . Na

SR CA

LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.

Double bond geometry as shown.




● Na


2 REFERENCES IN FILE CA (1967 TO DATE)
2 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 126:144046 Beta-lactam preparation. Harris, Michael Anthony; Saunders, Richard Neville (Pfizer Limited, UK). Brit. UK Pat. Appl. GB 2300856 A1 19961120, 15 pp. (English). CODEN: BAXXDU. APPLICATION: GB 1995-10126 19950516.

GI



I



II

AB Title compds. I [R = substituent; R1 = H, OMe, NHCHO; R2 = acyl; CO2R3 = CO2H, CO2-; R3 = protecting group; X = S, SO, SO2, O, CH2] are prepd. by base-induced cyclization of an azetidinone II [R4 = alkyl, aryl]. II are prepd. from the halide and P(OR4)3. Thus, 4-methoxybenzyl (2RS)-2-hydroxy-2-[(3R)(4R)-3-phenylacetamido-4-[(RS)-2-tetrahydrofuryl]carbonylmethylthio]azetidin-2-on-1-ylacetate was converted to the chloride and then to the phosphonate which was cyclized with NaH in PhMe to give 50% I [R = (RS)-2-tetrahydrofuryl, R1 = H, R2 = PhCH2CO, R3 = 4-MeC6H4CH2].

REFERENCE 2: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.


AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge,

Searched by: Mary Hale 308-4258 CM-1 1E01

carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L13 ANSWER 11 OF 13 REGISTRY COPYRIGHT 2002 ACS
 RN 141082-21-5 REGISTRY
 CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid,
 7-[[amino(4-hydroxyphenyl)acetyl]amino]-8-oxo-3-(tetrahydro-2-furanyl)-,
 monosodium salt, [6R-[3(S*),6.alpha.,7.beta.(R*)]]- (9CI) (CA INDEX NAME)
 FS STEREOSEARCH
 MF C19 H21 N3 O6 S . Na
 SR CA
 LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.




● Na

2 REFERENCES IN FILE CA (1967 TO DATE)
 2 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 126:144046 Beta-lactam preparation. Harris, Michael Anthony; Saunders, Richard Neville (Pfizer Limited, UK). Brit. UK Pat. Appl. GB 2300856 A1 19961120, 15 pp. (English). CODEN: BAXXDU. APPLICATION: GB 1995-10126 19950516.

GI



AB Title compds. I [R = substituent; R1 = H, OMe, NHCHO; R2 = acyl; CO2R3 = CO2H, CO2-; R3 = protecting group; X = S, SO, SO2, O, CH2] are prepd. by base-induced cyclization of an azetidinone II [R4 = alkyl, aryl]. II are prepd. from the halide and P(OR4)3. Thus, 4-methoxybenzyl (2RS)-2-hydroxy-2-[(3R)(4R)-3-phenylacetamido-4-[(RS)-2-

Searched by: Mary Hale 308-4258 CM-1 1E01

tetrahydrofuryl]carbonylmethylthio]azetidin-2-on-1-ylacetate was converted to the chloride and then to the phosphonate which was cyclized with NaH in PhMe to give 50% I [R = (RS)-2-tetrahydrofuryl, R1 = H, R2 = PhCH2CO, R3 = 4-MeC6H4CH2].

REFERENCE 2: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L13 ANSWER 12 OF 13 REGISTRY COPYRIGHT 2002 ACS

RN 141082-20-4 REGISTRY

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid,
7-[[[(5-amino-1,2,4-thiadiazol-3-yl)(methoxyimino)acetyl]amino]-8-oxo-3-(tetrahydro-2-furanyl)-, monosodium salt, [6R-[3(S*),6.alpha.,7.beta.(Z)]]-(9CI) (CA INDEX NAME)

OTHER CA INDEX NAMES:

CN 1,2,4-Thiadiazole, 5-thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid deriv.

FS STEREOSEARCH


MF C16 H18 N6 O6 S2 . Na

SR CA

LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.

Double bond geometry as shown.




● Na


2 REFERENCES IN FILE CA (1967 TO DATE)
2 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 126:144046 Beta-lactam preparation. Harris, Michael Anthony; Saunders, Richard Neville (Pfizer Limited, UK). Brit. UK Pat. Appl. GB 2300856 A1 19961120, 15 pp. (English). CODEN: BAXXDU. APPLICATION: GB 1995-10126 19950516.

GI



I



II

AB Title compds. I [R = substituent; R1 = H, OMe, NHCHO; R2 = acyl; CO2R3 = CO2H, CO2-; R3 = protecting group; X = S, SO, SO2, O, CH2] are prepd. by base-induced cyclization of an azetidinone II [R4 = alkyl, aryl]. II are prepd. from the halide and P(OR4)3. Thus, 4-methoxybenzyl (2RS)-2-hydroxy-2-[(3R)(4R)-3-phenylacetamido-4-[(RS)-2-tetrahydrofuryl]carbonylmethylthio]azetidin-2-on-1-ylacetate was converted to the chloride and then to the phosphonate which was cyclized with NaH in PhMe to give 50% I [R = (RS)-2-tetrahydrofuryl, R1 = H, R2 = PhCH2CO, R3 = 4-MeC6H4CH2].

REFERENCE 2: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge,

Searched by: Mary Hale 308-4258 CM-1 1E01

carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

L13 ANSWER 13 OF 13 REGISTRY COPYRIGHT 2002 ACS

RN 141082-16-8 REGISTRY

CN 5-Thia-1-azabicyclo[4.2.0]oct-2-ene-2-carboxylic acid, 7-[[[(2-amino-4-thiazolyl)(methoxyimino)acetyl]amino]-8-oxo-3-(tetrahydro-2-furanyl)-, monosodium salt, [6R-[6.alpha.,7.beta.(Z)]]- (9CI) (CA INDEX NAME)


FS STEREOSEARCH

MF C17 H19 N5 O6 S2 . Na .

SR CA

LC STN Files: CA, CAPLUS, TOXCENTER

Absolute stereochemistry.
Double bond geometry as shown.




● Na

3 REFERENCES IN FILE CA (1967 TO DATE)

3 REFERENCES IN FILE CAPLUS (1967 TO DATE)

REFERENCE 1: 126:144046 Beta-lactam preparation. Harris, Michael Anthony; Saunders, Richard Neville (Pfizer Limited, UK). Brit. UK Pat. Appl. GB 2300856 A1 19961120, 15 pp. (English). CODEN: BAXXDU. APPLICATION: GB 1995-10126 19950516.



GI



AB Title compds. I [R = substituent; R₁ = H, OMe, NHCHO; R₂ = acyl; CO₂R₃ = CO₂H, CO₂-; R₃ = protecting group; X = S, SO, SO₂, O, CH₂] are prepd. by base-induced cyclization of an azetidinone II [R₄ = alkyl, aryl]. II are prepd. from the halide and P(OR₄)₃. Thus, 4-methoxybenzyl (2RS)-2-hydroxy-2-[(3R)(4R)-3-phenylacetamido-4-[(RS)-2-tetrahydrofuryl]carbonylmethylthio]azetidin-2-on-1-ylacetate was converted to the chloride and then to the phosphonate which was cyclized with NaH in PhMe to give 50% I [R = (RS)-2-tetrahydrofuryl, R₁ = H, R₂ = PhCH₂CO, R₃ = 4-MeC₆H₄CH₂].

REFERENCE 2: 121:35060 Novel C-3 cyclic ether cephalosporins and their orally absorbed prodrug esters. Bateson, John H.; Burton, George; Fell, Stephen C. M.; Smulders, Hazel C. (Dep. Med. Chem., SmithKline Beecham Pharm., Betchworth/Surrey, RH3 7AJ, UK). J. Antibiot., 47(2), 253-6 (English) 1994. CODEN: JANTAJ. ISSN: 0021-8820.

GI



AB Cyclic ether cephalosporins I (R = Na, n = 1, 2) and their prodrug esters I (R = CH₂OCOCMe₃) were prepd. via Wittig cyclization of .beta.-lactam phosphoranes II. I (R = Na, n = 1, 2) were tested for bactericidal activity against several strains; I (R = Na, n = 1) was significantly more potent than I (R = Na, n = 2) and compared favorably with cefuroxime and cefetamet. The oral absorption of I (R = CH₂OCOCMe₃, n = 1, 2) was examd. in mice and compared with the .alpha.-acetoxyethyl ester of cefuroxime and the pivaloyloxymethyl ester of cefetamet.

REFERENCE 3: 116:255397 Preparation of 3-tetrahydrofurylcephem-3-

Searched by: Mary Hale 308-4258 CM-1 1E01

carboxylates and analogs as antibiotics. Bateson, John Hargreaves; Burton, George; Fell, Stephen Christopher Martin (Beecham Group PLC, UK). PCT Int. Appl. WO 9201696 A1 19920206, 147 pp. DESIGNATED STATES: W: AU, CA, CS, FI, HU, JP, KR, NO, PL, US; RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LU, NL, SE. (English). CODEN: PIXXD2. APPLICATION: WO 1991-GB1228 19910722. PRIORITY: GB 1990-16189 19900724; GB 1991-9540 19910502.

GI For diagram(s), see printed CA Issue.

AB Title compds. (I; R1 = H, MeO, HCONH; R2 = acyl; R3 = H, neg. charge, carboxy-protective group; R4 = .ltoreq.4 substituents selected from alkyl, alkenyl, OH, halo, alkoxy, etc.; X = O, CH2, SOn; n= 0-2; m = 1, 2) were prepd. Thus, Na 2-(2-tritylaminothiazol-4-yl)-2-(Z)-trityloxyiminoacetate was condensed with tert-butyl (6R, 7R)-7-amino-3-[(R)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylate to give, after deprotection, (6R, 7R)-7-[2-(2-aminothiazol-4-yl)-2-(Z)-hydroxyiminoacetamido]-3-[(RS)-tetrahydrofuran-2-yl]ceph-3-em-4-carboxylic acid which had MIC of 0.50 and 0.25 .mu.g/mL against Escherichia coli (NCTC 1048) and Staphylococcus aureus (Oxford), resp.

=> fil caol;s l13

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	529.63	529.84
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	-9.44	-9.44

FILE 'CAOLD' ENTERED AT 16:19:23 ON 18 JUN 2002
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.
 COPYRIGHT (C) 2002 AMERICAN CHEMICAL SOCIETY (ACS)

FILE COVERS 1907-1966
 FILE LAST UPDATED: 01 May 1997 (19970501/UP)

This file contains CAS Registry Numbers for easy and accurate substance identification. Title keywords, authors, patent assignees, and patent information, e.g., patent numbers, are now searchable from 1907-1966. TIFF images of CA abstracts printed between 1907-1966 are available in the PAGE display formats.

This file supports REGISTRY for direct browsing and searching of all substance data from the REGISTRY file. Enter HELP FIRST for more information.

L14 0 L13

=> log y

COST IN U.S. DOLLARS	SINCE FILE	TOTAL
	ENTRY	SESSION
FULL ESTIMATED COST	0.38	530.22
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE	TOTAL
	ENTRY	SESSION
CA SUBSCRIBER PRICE	0.00	-9.44

STN INTERNATIONAL LOGOFF AT 16:19:27 ON 18 JUN 2002

Searched by: Mary Hale 308-4258 CM-1 1E01